

22.07.2014

Emergency needs list for Thalassaemia patients in Gaza
(Period time 3 months)

N	Category	Details	Quantity	Patients number
1.	Exjade Drug “250mg”	This medication is used to treat iron overload caused by blood transfusions in adults and children at least 2 years old.	231 Packs	67
2.	Exjade Durg “500mg”	Used to treat chronic iron overload syndrome caused by a genetic blood disorder in adults and children who are at least 10 years old.	166 Packs	67
3.	Desferal (Deferoxamine) Drug 40-50mg/kg/day- patients	- This medication is used along with other treatments (such as causing vomiting with syrup of ipecac, stomach pumping) to treat sudden iron poisoning. It is most effective when given as soon as possible after the iron was eaten. -This medication can also be used to help get rid of iron in patients with high iron levels due to many blood transfusions.	28.200 vials	235
4.	Leucocytes Removal Filter Blood (Cat# F4C2491	Used to reduce complications associated with transfused white blood cells that are contained in units of red blood cells and platelets.	933 Filter	311
5.	Desferal pump	-iron chelation machine - used for getting rid of iron in our body due to blood transfusion	235	235
6.	Zoledroic Acid (Aclasta)	used to treat osteoporosis	85	85
7.	Somatropin – Nutropin (growth hormonal Medication)	- (1-8 year)2 injection per day	1300 injection	7
		- (12-16 year) 3 injection per day	19,710 injection	73